

The pair production of charmed mesons in the photon-photon interaction.

A.V. Berezhnuy, V.V. Kiselev, A.K.Likhoded

Abstract

In the framework of the constituent quark model the exclusive pair production of the charmed mesons in the photon-photon interaction is calculated. The comparison of these predictions with the ones of the heavy quark effective theory is performed. It is shown, that the light valence quark of D-meson plays the essential role not only in the hadronization process, but also in the production of the heavy c-quark. Moreover, it is shown, that because of the strong interaction between the initial photon field and the light quark charge the similar situation is kept even in the limit $m_Q \rightarrow \infty$, whence it is concluded, that the application of the heavy quark effective theory in the case of the photon-photon interaction is not valid because it does not take into account one of the dominant mechanisms of the heavy meson production.

Introduction

Recent years the big volume of work devoted to the experimental study of the photon-photon interactions, in particular, to the study of the photonic production of the charmed particles was carried out at the accelerator LEP. In this connection, and also in connection with prospects for construction of the largest e^+e^- -collider TESLA (DESY) there is a necessity of the detailed discussion of the photon-photon interaction mechanisms at large energies. In the given work we shall consider some questions concerning the photonic production of heavy quarks, namely, we shall discuss features of the exclusive and the inclusive charmed meson pair production and estimate its contribution into the total cross section of the charm production. In the framework of the constituent quark model (CQM) we shall show, that the light valence quark inside D-meson plays the essential role not only in the hadronization process, but also in the hard production of the heavy quark. Because of strong interaction of

the photonic field with the light quark charge the similar situation is kept even in a limit $m_Q \rightarrow \infty$.

It is worth to notice, that for the exclusive pair production of the charmed meson in the e^+e^- -annihilation CQM and the heavy quark effective theory (HQET) give the same prediction for $m_Q \rightarrow \infty$. Nevertheless at the conventional values of masses of the light and heavy quarks the results of CQM for the e^+e^- -interactions strongly differ from the results of the HQET.

1 The pair production of the charmed mesons in the e^+e^- -annihilation.

We begin our article with the discussion of the heavy meson production in the e^+e^- -annihilation, because it is the most simple process, and because it is considered in detail in the literature.

The work[1] presents one of the first attempts to formulate the HQET for the case of the D -meson production in the e^+e^- -annihilation. The neglect of spin-spin interaction between the heavy and light quarks, which is suppressed as $1/m_Q$, has allowed to obtain the interesting ratio for the pair production of D and D^* -meson near the threshold (see the Appendix):

$$\sigma_{D\bar{D}} : \sigma_{D\bar{D}^*+D^*\bar{D}} : \sigma_{D^*\bar{D}^*} = 1 : 4 : 7 \quad (1)$$

More detailed analysis of the exclusive $D\bar{D}$ -pair production in the e^+e^- -interaction on the basis of the CQM, was carried out in work [2]. It is worth to remind, that CQM is based on the assumptions, that in the partonic distributions of the final meson there are the valence quark contributions, and that both valence quarks are produced in the hard process followed by the fusion of these quarks into the meson.

The analytical expressions for the exclusive meson pair production look as follows [2]:

$$\begin{aligned} \sigma(e^+e^- \rightarrow (Q\bar{q})_P(\bar{Q}q)_P) &= \frac{\pi^3\alpha_s^2(4m_q^2)\alpha_{em}^2}{3^7 4m_q^6} \frac{m_Q^2}{M^2} f_P^4 (1-v^2)v^3 \times \\ &\times \left\{ 3e_Q \left(\frac{2m_q}{m_Q} - 1 + v^2 \right) - 3e_q \left[2 - (1-v^2) \frac{m_q}{m_Q} \right] \frac{m_q^3\alpha_s(m_Q^2)}{m_Q^3\alpha_s(4m_q^2)} \right\}^2, \end{aligned} \quad (2)$$

$$\sigma(e^+e^- \rightarrow (Q\bar{q})_P(\bar{Q}q)_V) = \frac{\pi^3\alpha_s^2(4m_q^2)\alpha_{em}^2}{3^7 4m_q^6} \frac{m_Q^2}{M^2} f_P^2 f_V^2 (1-v^2)v^3 \left[3e_Q + 3e_q \frac{m_q^3\alpha_s(m_Q^2)}{m_Q^3\alpha_s(4m_q^2)} \right]^2, \quad (3)$$

$$\sigma(e^+e^- \rightarrow (Q\bar{q})_V(\bar{Q}q)_V) = \frac{\pi^3 \alpha_s^2 (4m_q^2) \alpha_{em}^2}{3^7 4m_q^6} f_V^4 (1-v^2) v^3 \left[3e_Q - 3e_q \frac{m_q^3 \alpha_s (m_Q^2)}{m_Q^3 \alpha_s (4m_q^2)} \right]^2 \times$$

$$\times \left[3(1-v^2) + (1+v^2)(1-a)^2 + \frac{a^2}{2}(1-v^2)(1-3v^2) \right], \quad (4)$$

where m_q and m_Q are masses of the light and heavy quarks correspondingly; f_P and f_V are leptonic coupling constants for the pseudoscalar and vector meson states correspondingly; $M = m_q + m_Q$; $v = \sqrt{1 - 4M^2/s}$; and the parameter a is expressed by the formula:

$$a = \frac{m_Q}{M} \frac{1 - \frac{e_q}{e_Q} \frac{m_q^4}{m_Q^4} \frac{\alpha_s(4m_Q^2)}{\alpha_s(4m_q^2)}}{1 - \frac{e_q}{e_Q} \frac{m_q^3}{m_Q^3} \frac{\alpha_s(4m_Q^2)}{\alpha_s(4m_q^2)}}.$$

At $m_Q \rightarrow \infty$ the ratio of the cross sections on the threshold looks like:

$$\sigma_{PP} : \sigma_{PV} : \sigma_{VV} = 1 : 4 \frac{f_V^2}{f_P^2} : 7 \frac{f_V^4}{f_P^4}. \quad (5)$$

Thus if one neglects the difference between the values of f_V and f_P , then (5) coincides with (1).

The heavy quark effective theory predicts the ratio which at a threshold also coincides with (1):

$$(1+h) : \frac{s}{M^2} : 3 \left(1 + \frac{s}{3M} + h \right),$$

where the α_s -correction has the following form:

$$h = -\frac{2\alpha_s}{3\pi} \sqrt{1 - \frac{4M^2}{s}} \ln \left(\frac{s}{2M^2} - 1 + \frac{s}{2M^2} \sqrt{1 - \frac{4M^2}{s}} \right).$$

The four Feynmann diagrams of $O(\alpha_s^2 \alpha_{em}^2)$ order contribute to the cross sections (2), (3), (4) (see Fig.1). These diagrams can be separated into two gauge invariant pairs. The first pair (two top diagrams of Fig.1) corresponds to the interaction of the virtual photon with the heavy quark and is proportional to the heavy quark charge squared. The value of α_s for this diagram pair is calculated at the scale of $4m_q^2$, because the gluonic propagator couples with the two light quarks. The second




Figure 1: The diagrams of the meson production in the e^+e^- -annihilation.

diagram pair (the two bottom diagrams of the Fig. 1) corresponds to the interaction of virtual photon with the light quark and is proportional to the light quark charge. For these diagrams α_s is calculated at the scale $4m_Q^2$. It is clear, that this contribution is suppressed by the factor m_q^2/m_Q^2 due to the gluonic propagator coupled with the pair of heavy quarks.

One can see from (2), (3), (4), that at $m_Q \rightarrow \infty$ it is possible to neglect the interaction of the light quark with the initial virtual photon. It is necessary to note, that the contribution of that interaction is small already for the masses of constituents of the D -meson (for present calculations the following values of masses were chosen: $m_c = 1.8$ GeV and $m_q = 0.2$ GeV). However, the spin-spin interaction at these values of masses plays the essential role and brakes the ratio (1):

$$\sigma_{D\bar{D}} : \sigma_{D\bar{D}^*+D^*\bar{D}} : \sigma_{D^*\bar{D}^*} \approx 1 : 8 : 14.$$

So, the $D\bar{D}$ -pair production in the e^+e^- -annihilation occurs as follows: a pair of heavy quarks is produced in a hard process, and then these quarks are hadronized by means of interaction with the light quark sea. It is worth to remind, that in the inclusive production of D -mesons in the e^+e^- -annihilation, where the fragmentation mechanism can be applied[3], the c -quarks are hadronized by the same manner. Thus, the role of light quark in the hard process of the c -quark production in the e^+e^- -annihilation is insignificant. On the contrary, in the photon-photon production the light quark role is essential as will be shown bellow.

2 The exclusive $D\bar{D}$ -pair production in the photon-photon interaction

In the framework of the CQM the photonic exclusive production of two charmed mesons is described by twenty Feynmann tree level diagrams (see the Fig. 2), which can be separated into three gauge invariant groups. The first of them corresponds to a case when the heavy quark radiate the gluon which splits into a pair of light quarks (diagrams 1-6 of the Fig. 2). The contribution of this group is proportional to the square of the heavy quark charge. In the second group of diagrams (the diagram 7-12), on the contrary, the heavy quark production occurs by means of the splitting of the gluon which is radiated by the light quark. These diagrams are proportional to the square of the light quark charge and suppressed by the factor m_q^2/m_Q^2 due to the gluonic propagator which is proportional to $1/m_Q^2$.

In the third group of diagrams (the diagram of 13-20 Fig. 2) each pair of quarks is connected to its "own" γ -quantum. Our calculations show, that the interaction of the




Figure 2: The diagrams of the photonic production of two heavy mesons.

light quark with the initial photon field described by these diagrams is not suppressed and plays the essential role in the exclusive pair production of the charmed meson in the photon-photon interaction. This interaction leads to the discrepancy between the predictions of HQET and CQM, which is not negligible even at $m_Q \rightarrow \infty$. Moreover, the interaction of the light constituent quark with the initial photon is so large, that HQET can not be applied in this case.

Nevertheless, if one puts the light quark charge equal to zero in our calculations (this case corresponds to the rejection of the diagrams of the second and third groups, i.e. 7-20 of Fig. 2) then the conformity between CQM and HQET is restored. Thus, at zero charge of the light quark and $m_Q \rightarrow \infty$ we have received on the threshold the following ratio:

$$\sigma_{PP} : \sigma_{PV} : \sigma_{VV} = 1 : 0 : 3, \quad (6)$$

predicted in the work[4], where the approach [1] is applied to the case of photonic production (see Appendix). If the light quark charge is differ from zero then its interaction with the photonic field becomes essential for the $D\bar{D}$ -pair production. Moreover, our calculation show, that $m_Q \rightarrow \infty$ and $2m_Q/s \rightarrow 1$ lead to $\sigma_{VV}/\sigma_{PP} \rightarrow \infty$. It means that the mechanism of the exclusive production based on the production of the heavy quark pair followed by their hadronization appears not valid. As it is shown in the work [5], such approach is not valid in the case of the inclusive production too.

In the Fig. 3 we present the exclusive cross sections of the photonic pair production of the charmed mesons σ_{PP} (open circles), σ_{PV} (open triangles), σ_{VV} (open squares) calculated by us in the framework of the CQM as a function of $\Delta = \sqrt{s} - \sqrt{s_{th}}$ with the following values of parameters:

$$m_c = 1.5 \text{ GeV};$$

$$m_q = 0.3 \text{ GeV};$$


$$\alpha_s = 0.3.$$

As well as in the case of production in the e^+e^- -annihilation, the cross sections σ_{PP} , σ_{PV} and σ_{VV} are proportional to f_P^4 , $f_P^2 f_V^2$ and f_V^4 , accordingly.

The values of f_P and f_V estimated in the frame work of the sum rules, as well as in the frame work of the lattice calculations are about $f_P \sim f_V \sim 200 - 300 \text{ MeV}$. However, the absolute normalization of the cross sections in our model depends not only on these constants, but also on the light quark mass. The variation of the light quark mass can result in the change of the cross section normalization.

On the other hand, we can normalize the values of f_P and f_V on the base of the fragmentation probability $c \rightarrow D^*$ in the e^+e^- -interaction at large energies ($W(c \rightarrow$

$\sigma, \text{ nb}$



$\Delta, \text{ GeV}$

Figure 3: The exclusive cross section of the photonic pair production of the charged charmed mesons σ_{PP} (the open circles), σ_{PV} (the open triangles), σ_{VV} (the open squares) as a function of $\Delta = \sqrt{s} - \sqrt{s_{th}}$. The black markers show the same dependences at a zero charge of the light quark.

$D^*) = 0.22$). Thus the values of m_q and $f_{P,V}$ appear correlated to each other, that allows to minimize the cross section uncertainty within 2-2.5. However, with such normalization the constant $f_{P,V}$ appears overestimated at all reasonable values of the light quark mass. We believe, that this overestimate of the constant $f_{P,V}$ is due to the account for the contribution of the excited state decays, for example, of $D^{**} \rightarrow D^* + X$. In the exclusive production such situation should not be observed. Thus, the question of the absolute normalization for the exclusive channels remains open and the values in Fig.3 should be considered as the top limit of the cross section. For all conclusions connected to the relative yield of D -mesons the discussed circumstances are not essential.

Let us notice, that near the threshold $\Delta \approx |k|^2/M$, where $|k|$ is the module of three-momentum of the final meson in the center-of-mass frame of the charmed meson pair. The twice logarithmic scale of the picture reveals, that the cross section depends on Δ near the threshold as:

$$\sigma_{VV}, \sigma_{PP} \sim \Delta^{1/2} \sim |k|,$$

$$\sigma_{PV} \sim \Delta^{3/2} \sim |k|^3.$$

The point is that in the $\gamma\gamma$ -interaction the $D\bar{D}$ - and $D^*\bar{D}^*$ -pairs are produced in S-wave state (L=0), and $D\bar{D}^*$ -pairs are produced in P-wave state (L=1). It is known, that near the threshold the behavior of reactions with two particles in the final state submits to the law:

$$\sigma \sim |k|^{2L+1}. \quad (7)$$

The diagrams of Fig.2 which describes the processes $\gamma\gamma \rightarrow D\bar{D}$, $\gamma\gamma \rightarrow D^*\bar{D}^*$, $\gamma\gamma \rightarrow D\bar{D}^*$ are written at the quark level and the "twisting" of the final mesons into the necessary wave occurs automatically. Hence, the observable behavior of the cross sections near threshold is the additional confirmation of the calculation correctness.

It is apparent from Fig. 3, that the HQET ratio (6) is broken. Our calculations show, that near the threshold

$$\sigma_{PP} : \sigma_{PV} : \sigma_{VV} \approx 1 : 0 : 13.4.$$

It is interesting to consider the case when the interaction of the light quark with the initial photon can be neglected i.e. when the electric charge of the light quark is put equal to zero. The black markers in Fig. 3 show the results of the calculations of the charmed meson pair production cross sections at a zero light quark charge.

One can see, that near the threshold the ratio $\sigma_{PP} : \sigma_{PV} : \sigma_{VV}$ is practically the same as predicted by (6). It is interesting to mention, that the contribution from the interaction between the photon and the light quark can be positive (the case of $D^*\bar{D}^*$ -pair production), as well as negative (the case of $D\bar{D}$ or $D^*\bar{D}$ -pair production). Let us note also, that contribution of this interaction has the minimal absolute value for the $D^*\bar{D}$ -pair production, i.e. when the mesons are formed in the P -wave state.


The extremely essential role of the light quark in the charmed meson production can be understood from the difference between the production of neutral mesons and the charged ones. This difference can be clearly seen in Fig. 4, where the Δ -dependence of the neutral meson cross section (the black markers) is shown in comparison with the same dependence for charged mesons (the open ones). For example, near the threshold for the charged mesons $\sigma_{D^+D^-} : \sigma_{D^{*+}D^{*-}} \approx 1 : 13.4$ in the contrary to $\sigma_{D^0\bar{D}^0} : \sigma_{D^{*0}\bar{D}^{*0}} \approx 1 : 2.7$ for the neutral ones.

It is necessary to note, that the least difference between the charged and neutral meson production is observed for the production of $D\bar{D}^*$ -pairs i.e. when the mesons are produced in the P -wave state.

The more striking differences can be observed between the neutral and charged B-meson production. In Fig. 5 the production cross section of the neutral B-meson is given as a function of $\Delta = \sqrt{s} - \sqrt{s_{th}}$ for $m_b = 5$ GeV and $m_q = 0.2$. In the case of D-meson production the cross section dependences on Δ have the qualitatively identical form both for the charged particles and for the neutral ones (see Fig. 4). For the B-meson production it is not so. One can see in Fig. 5, that at $\Delta \approx 0.12$ GeV ($|k| \approx 0.8$ GeV) the interferential minimum is observed for the production of two charged pseudo-scalar mesons B^+ and B^- and not observed for the pair production of the neutral pseudo-scalar mesons B^0 and $\bar{B}^0$.

So, for all interaction energies it is necessary to take into account the light quark contribution into the photonic production of heavy mesons. The models which are not taking into account this contribution can not be applied even for the qualitative description of the considered process. It is clear from the analysis of the photonic pair production of the heavy mesons, that the naive ideas which transferred from the e^+e^- -annihilation appear not valid. For the processes $\gamma\gamma \rightarrow D\bar{D}$ and $\gamma\gamma \rightarrow B\bar{B}$ the role of light quarks appears essential both in the soft process of the hadronization and in the hard pair production of the heavy quarks. Unfortunately, the experimental data on the heavy meson pair production near threshold do not exist yet. Such data could throw light on the dynamics of production of mesons with the heavy quarks.


σ , nb



Δ , GeV

Figure 4: The exclusive cross section of the photonic pair production for the charged charmed mesons σ_{PP} (open circles), σ_{PV} (open triangles), σ_{VV} (open squares) as function of Δ in comparison with the same dependences for the neutral mesons (black markers).

σ , nb



Δ , GeV

Figure 5: The photonic production of the charged B -mesons (black markers) and the neutral ones (open markers) as a function of Δ .

3 The inclusive heavy meson production in the photon-photon interaction

As it was already mentioned, the exclusive heavy meson production data would allow to compare the predictions of various models, however the opportunities of experiment in this direction of researches are strongly limited. For today the majority of measurements are made for the inclusive D -meson production. To describe this process a number of the calculations is carried out in the framework of NLO QCD for the massive c -quarks [7], as well as for c -quarks [8]. These calculations result in the satisfactory description of the data, however, the use of the fragmentation mechanism result in specific predictions for the relative yield of charged and neutral mesons. In these models the D^* -yield does not depend on the meson type: $\sigma_{D^{*0}} = \sigma_{D^{*+}}$.

Let us consider now the prediction of our model for the inclusive heavy meson production cross section. The diagrams we need are analogous to the ones of Fig. 2. The only difference is that the second pair of heavy and light quarks is not fused in the meson. Thus we take into account the resonant part of the spectrum for one quark pair and the continuum for another. To calculate the inclusive production cross section we use the values of parameters which describes the photoproduction of the D^* -mesons on the accelerator HERA [5], as well as in our calculation of the exclusive charmed meson production. To estimate the inclusive production cross section we shall take into account the octet contribution which is absent in the exclusive production. In the case of the photonic inclusive production of D -mesons the octet contribution differs from the singlet one only in the general coefficient. If one use the parameters, from [5], then the octet contribution amounts 16% of the total cross section of the D -meson production.

One can see from Fig. 6 that our predictions for the inclusive production describe L3 experiment data in the order of magnitude (open circles). As our calculations show, the nonzero light quark charge results in the value of ratio $\sigma_{D^{*+}}/\sigma_{D^{*0}}$ differ from unit. This is valid for the inclusive production too: our calculations show, that the inclusive D^{*0} -meson yield approximately twice surpasses the yield of the D^{*+} -meson, while the fragmentation model predicts $\sigma_{D^{*+}}/\sigma_{D^{*0}} = 1$.

At energies larger than 15 GeV (the range where $c\bar{q}$ -pair produced mainly at large invariant masses) the total cross section calculated in our model begins to decrease and underestimate the experimental data. In this energy range it is not enough to consider the resonant contribution (open triangles), and it is necessary to take into account the contribution of continuum (open squares). To calculate the contribution of the continuum we do not fuse now the light quarks with the c -quarks: $\gamma\gamma \rightarrow c\bar{c}q\bar{q}$. In the conception of structure functions the considered mechanism means the account

σ , nb




Figure 6: The charm production cross section $\gamma\gamma \rightarrow c\bar{c} + X$ (open circles) in comparison with the L3 Collaboration experimental data[6].

of the hadronic structures of the photon. Our approach allow to take into account only a part of the photon structure. That is why we underestimate the production cross section, and the additional part of the photon structure function is necessary to total value of the cross section. We expect for the contribution of continuum, that the yields of the charged mesons and neutral ones should be identical. Thus the ratio $\sigma_{D^{*+}}/\sigma_{D^{*0}}$ could serve as a good indication of the production mechanism. The calculations which have been carried out in the works [7, 8] show that the use of the total structure function allows to describe experimental data well.

Conclusion

In the framework of CQM we have analysed the exclusive production of the heavy mesons in the $\gamma\gamma$ -collisions near the threshold, as well as the inclusive heavy meson production at high energies. Near the threshold we have calculated the relative contribution of the different meson pairs and have found out that the calculation results essentially depends on the electric charge of the light constituent quark. For zero electric charge of the light quark we obtain, that the production process looks

like the heavy quark production followed hadronization according to HQET. At the real values of charges the strong deviation from the predictions of the HQET takes place. Thus, our analysis of the photon-photon production of the heavy meson pairs shows, that the naive conception which is transferred from the e^+e^- -annihilation appear not valid. In the processes $\gamma\gamma \rightarrow D\bar{D}$ and $\gamma\gamma \rightarrow B\bar{B}$ the role of the light quarks is essential both in the soft hadronization process and in the hard process of the heavy quark pair production. Unfortunately, in the framework of our model it is impossible to use the same values of constants f_P and f_V to calculate both the exclusive channel and the inclusive one. The use of the values of f_P and f_V , which describes process $\gamma p \rightarrow D^* + X$, leads to the large overestimate of the production cross section near the threshold. We suppose, that such overestimation takes place because the values f_P and f_V effectively account for the contributions from the decays of the excited states, for example, from the decay $D^{**} \rightarrow D^* + X$. So, these values can not be applied to the case of the exclusive meson production. Nevertheless, our model gives reasonable estimation of the relative contributions into the cross section of the different heavy meson pairs ($D^0\bar{D}^0$, $D^{0*}\bar{D}^0$, $\bar{D}^0D^{0*}$, $D^{0*}\bar{D}^{0*}$, D^+D^- , $D^+\bar{D}^{*-}$, D^-D^{+*} , $D^{+*}D^{*-}$), as well as reasonable predictions of the dependence of the cross section on the interaction energy and the shape of the differential distributions.

Also in the framework of the CQM we calculated the cross section of the inclusive charmed meson production in the photon-photon interaction. The cross section prediction of the inclusive production calculated in the frame work of CQM is in accordance with the experiment data at low energies of the $\gamma\gamma$ -interactions if one choose the parameter values which describes the process $\gamma p \rightarrow D^* + X$ at HERA [5]. It is not unexpected for us because the contribution of the low γg -interaction energies is the most essential for the D^* -meson photoproduction at the accelerator HERA . The continuum contribution should increase as a function of the interaction energy. Thus at large energies the character of the calculations should be the same, as in the works [7, 8], which explain the cross section increasing for the process $\gamma\gamma \rightarrow c\bar{c}$ with the help of the account for the contribution of the light constituents in the initial photon structure.

The work is supported in part by the grants of RFBR 99-02-6558 and 00-15-96645, the grant of the Education Ministry of Russian Federation E00-33-062 and the grant CRDF MO-001-0.

Appendix

In the given appendix we show how to derive the ratio $\sigma_{PP} : \sigma_{PV} : \sigma_{VV}$ near the threshold for the heavy meson production in the e^+e^- -annihilation, as well as in the

$\gamma\gamma$ - interaction in the approach of the heavy quark effective theory.

Let us accept the following designations:

$Q_{+(-)}$ ($\bar{Q}_{+(-)}$) is the wave function of the heavy quark (antiquark) with the spin projection $+1/2(-1/2)$ on the chosen axis;

$q_{+(-)}$ ($\bar{q}_{+(-)}$) is the wave functions of the light quark (antiquark).

In the framework of HQET the heavy quarks are produced in the e^+e^- -annihilation the by means of the virtual photon decay, so, these quarks are in $J^P = 1^-$ state. The spin part of the wave function of such state can be expressed through the spinor wave functions of the quarks as follows:

$$\hat{\Psi}_{Q\bar{Q}} = (\Psi_{Q\bar{Q}}^{+1}, \Psi_{Q\bar{Q}}^0, \Psi_{Q\bar{Q}}^{-1}) = (Q_+\bar{Q}_+, \frac{Q_+\bar{Q}_- + Q_-\bar{Q}_+}{\sqrt{2}}, Q_-\bar{Q}_-).$$

Thus the light quarks should be in the 0^+ -state. This state can be obtained in the P-wave as follows:

$$\psi_{q\bar{q}} = (n_{-1}\Psi_{q\bar{q}}^{+1} + n_0\Psi_{q\bar{q}}^0 + n_{+1}\Psi_{q\bar{q}}^{-1})/\sqrt{3},$$

where $\hat{n}$ is the orbital part of the wave function of the light quark pair, and $\hat{\Psi}_{q\bar{q}}$ is the spin part of the one:

$$\hat{\Psi}_{q\bar{q}} = (\Psi_{q\bar{q}}^{+1}, \Psi_{q\bar{q}}^0, \Psi_{q\bar{q}}^{-1}) = (q_+\bar{q}_+, \frac{q_+\bar{q}_- + q_-\bar{q}_+}{\sqrt{2}}, q_-\bar{q}_-).$$

So, the total wave function looks like follows:

$$\hat{\Psi} = \hat{\Psi}_{Q\bar{Q}} \cdot (n_{-1}\Psi_{q\bar{q}}^{+1} + n_0\Psi_{q\bar{q}}^0 + n_{+1}\Psi_{q\bar{q}}^{-1})/\sqrt{3}. \quad (8)$$

The right part of equation (8) includes the products of the spinor wave functions of the quarks $Q_{+(-)}\bar{q}_{+(-)}$ and $\bar{Q}_{+(-)}q_{+(-)}$, which can be expressed through the spin functions of the final mesons:

$$P = \frac{Q_+\bar{q}_- - Q_-\bar{q}_+}{\sqrt{2}}, \quad (9)$$

where P is the wave function of the pseudoscalar meson;

$$V_{+1} = Q_+\bar{q}_+, \quad (10)$$

$$V_0 = \frac{Q_+\bar{q}_- + Q_-\bar{q}_+}{\sqrt{2}}, \quad (11)$$

$$V_{-1} = Q_{-}\bar{q}_{-}, \quad (12)$$

where P and V are the wave functions of the pseudoscalar and vector mesons, correspondingly.

One can conclude from (9) and (11) that

$$Q_{+}\bar{q}_{-} = \frac{V_0 + P}{\sqrt{2}}, \quad (13)$$

$$Q_{-}\bar{q}_{+} = \frac{V_0 - P}{\sqrt{2}}. \quad (14)$$

From the equations (10,12,13,14) the following expressions for the components of the total wave function (8) can be derived :

$$\begin{aligned} \Psi^{+1} = & \frac{1}{2\sqrt{3}} \left(n_{+1}P\bar{P} + \{n_{+1}P\bar{V}_0 + n_{+1}V_0\bar{P} + n_0P\bar{V}_{+1} + n_0V_{+1}\bar{P}\} + \right. \\ & \left. + [n_{+1}V_0\bar{V}_0 + n_0V_0\bar{V}_{+1} + n_0V_{+1}\bar{V}_0 + 2n_{-1}V_{+1}\bar{V}_{+1}] \right), \end{aligned} \quad (15)$$

$$\begin{aligned} \Psi^0 = & \frac{1}{2\sqrt{3}} \left(-n_0P\bar{P} + \{n_{+1}P\bar{V}_{-1} + n_{+1}V_{-1}\bar{P} - n_{-1}P\bar{V}_{+1} - n_{-1}V_{+1}\bar{P}\} + \right. \\ & \left. + [n_{+1}V_0\bar{V}_{-1} + n_{+1}V_{-1}\bar{V}_0 + n_0V_0\bar{V}_0 + n_0V_{+1}\bar{V}_{-1} + n_0V_{-1}\bar{V}_{+1} + n_{-1}V_{+1}\bar{V}_0 + n_{-1}V_0\bar{V}_{+1}] \right), \end{aligned} \quad (16)$$

$$\begin{aligned} \Psi^{-1} = & \frac{1}{2\sqrt{3}} \left(n_{-1}P\bar{P} - \{n_{-1}P\bar{V}_0 + n_{-1}V_0\bar{P} + n_0P\bar{V}_{-1} + n_0V_{-1}\bar{P}\} + \right. \\ & \left. + [n_{-1}V_0\bar{V}_0 + n_0V_0\bar{V}_{-1} + n_0V_{-1}\bar{V}_0 + 2n_{+1}V_{-1}\bar{V}_{-1}] \right). \end{aligned} \quad (17)$$

As we believe, that the spins of the heavy quark and the light quark do not interact with each other and with the orbital moment, the all term in the right part of the equations (15,16,17) give the identical contributions to the square of the total wave function $\hat{\Psi}$. The result (5) becomes obvious, because all these terms are orthogonal to each other.

In the case of $\gamma\gamma$ -interactions two initial photons can be in the following states: 0^+ , 0^- , 2^+ . And the 0^+ -state can be produced in the S -wave only. Since at the threshold survive contributions of the state with a minimal wave number (see (7)), we shall consider 0^+ . Such state can be made of two pseudo-scalar functions:

$$\Psi_{Q\bar{Q}} = \frac{Q_{+}\bar{Q}_{-} - Q_{-}\bar{Q}_{+}}{\sqrt{2}}$$

and

$$\Psi_{q\bar{q}} = \frac{q_+\bar{q}_- - q_-\bar{q}_+}{\sqrt{2}}$$

by the following manner:

$$\Psi = \Psi_{Q\bar{Q}} \cdot \Psi_{q\bar{q}}. \quad (18)$$

Using (10,12,13,14) to transform the equation (18), we obtain:

$$\Psi = \frac{1}{2}(-P\bar{P} + V_0\bar{V}_0 - V_{-1}\bar{V}_{+1} - V_{+1}\bar{V}_{-1}),$$

whence the ratio for the photonic production near the threshold follows:

$$\sigma_{PP} : \sigma_{PV} : \sigma_{VV} = 1 : 0 : 3$$

References

- [1] A. De Rújula, Howard Georgi and S. L. Glashow, *Phys. Rev. Lett.* **37**, 398 (1976).
- [2] V. V. Kiselev, *International Journal of Modern Physics* **A10**, 465 (1995).
- [3] V.G. Kartvelishvili, A.K. Likhoded and V.A. Petrov, *Phys.Lett.* **B78**, 615 (1978).
- [4] V.G. Kartvelishvili and A.K. Likhoded *JETP Lett.* **26**, 54 (1976).
- [5] A. V. Berezhnoy, V. V. Kiselev and A. K. Likhoded, *Phys.Rev.* **D62**, 074013 (2000).
- [6] M. Acciarri et al. (L3 Collab.), *Phys.Lett.* **B514**, 19 (2001).
- [7] S. Frixione, M. Krämer and Laenen, *Nucl.Phys.* **B571**, 169 (2000); *J.Phys.* **G26**, 723 (2000).
- [8] J. Binnewies, B.A. Kniehl and G. Kramer, *Phys.Rev.* **D53**, 6110 (1996); *Phys.Rev.* **D58**, 014014 (1998).